

AD-A115 945

COLUMBIA UNIV NEW YORK DEPT OF CHEMISTRY
INDUCTION OF OLEFIN METATHESIS BY PHENYLACETYLENE PLUS TUNGSTEN--ETC(U)
JUN 82 T J KATZ, C HAN

F/6 11/9

N00014-79-C-0683


NL

UNCLASSIFIED

TR-4

1 OF 1
404
F-144

END
DATE
FILED
07-32
OTC



AD A 11 5945

12

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

DINE FILE COPY

DD FORM 1 JAN 73 1473 EDITION OF 1 NOV 65 IS OBSOLETE

Accession For	
NTIS GRA&I	<input type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Special
A	

OFFICE OF NAVAL RESEARCH

Contract N00014-79-C-0683

Task No. NR 356-726

TECHNICAL REPORT NO. 4



Induction of Olefin Metathesis by Phenylacetylene

Plus Tungsten Hexachloride

by

Thomas J. Katz, Chien-Chung Han

Prepared for Publication

in

Organometallics

Columbia University
Department of Chemistry
New York, New York

June 21, 1982


Reproduction in whole or in part is permitted for
any purpose of the United States Government

This document has been approved for public release
and sale; its distribution is unlimited

The discovery by Masuda *et al.* that tungsten hexachloride initiates polymerization of phenylacetylene¹ suggests the following proposition: that phenylacetylene may effectively replace the highly reactive organo-aluminum component in the mixture with tungsten hexachloride that is the archetypical² and still standard initiator for olefin metathesis.³ For if the metal-catalyzed polymerization of acetylenes is an olefin metathesis,⁴⁻⁶ a metallic compound that induces an acetylene to polymerize should itself be induced by the acetylene to metathesize olefins. The idea was demonstrated recently with phenylacetylene plus (phenylmethoxycarbene)pentacarbonyltungsten.⁶ It is shown here also for phenylacetylene plus tungsten hexachloride, a combination that is more available, acts faster, induces fair stereoselectivity, and can be used unprotected in the atmosphere.⁷

Fig 1

Figure 1 shows phenylacetylene's effect on the metathesis of cis-2-pentene¹⁴ according to eq 1.¹⁵ As phenylacetylene is added, the



amount of reaction, negligible in its absence, rises dramatically and then falls. Complete reaction is easily achieved, but only when the amount of acetylene is appropriate.

Table I

Table I similarly shows phenylacetylene's ability to stimulate tungsten hexachloride to polymerize three cyclic olefins (eq 2), reactions that (as seen in the table) occur inappreciably, if at all, when





Figure 1. The extent of reaction according to eq 1 as a function of x . Percent metathesis is recorded as $200r/(1 + 2r)$, where r is the ratio in moles of either 2-butene to 2-pentene or 3-hexene to 2-pentene. The bars span the two values associated with r measured in the two ways.



phenylacetylene is absent. That the polymers formed are indeed polyalkenamers, not saturated vinyl polymers, is attested to by their ^1H NMR spectra displaying only the required resonances and intensities. The latter measure contamination by saturated hydrocarbon amounting to no more than 4-9% ($\pm 4\%$).

Table II

Table II shows that for cyclopentene, as for 2-pentene, while small amounts of phenylacetylene induce metathesis, larger amounts quench reaction. The decreased molecular weights (also recorded there) when the amount of acetylene is large imply that growing chains rather than initiating centers are being extinguished. This termination may involve eq 3, the transformation of a simpler metal-carbene into one more highly substituted and with conjugating groups.^{6b} One might speculate that the initiation involves eq 4, possibly accompanied by



reaction with additional acetylene reducing the tungsten's oxidation state.¹⁸

A hypothesis associating diminished stereoselectivity with the presence of Lewis acids^{3e,19} suggests that stereoselectivity should increase when an aluminum halide is replaced by phenylacetylene and decrease when (phenylmethoxycarbene)pentacarbonyltungsten is replaced by tungsten hexachloride. The stereoselectivities recorded in Table II and the stereochemistries of the 2-butenes (initially 69.6% cis) and 3-hexenes (initially ca. 54% cis) in Figure 1 accord with this hypothesis.²⁰

Acknowledgments. We are grateful to the National Science Foundation (CHE-81-08998) and the U.S. Office of Naval Research for support.

References and Notes

- (1) Masuda, T.; Hasegawa, K.; Higashimura, T. Macromolecules 1974, 7, 728.
- (2) Natta, G.; Dall'Asta, G.; Mazzanti, G. Angew. Chem. Int. Ed. Engl. 1964, 3, 723.
- (3) (a) Banks, R.L. Catalysis (London) 1981, 4, 100. (b) Calderon, N.; Lawrence, J. P.; Ofstead, E. Adv. Organomet. Chem. 1979, 17, 449. (c) Grubbs, R. H. Prog. Inorg. Chem. 1978, 24, 1. (d) Rooney, J. J.; Stewart, A. Catalysis (London) 1977, 1, 277. (e) Katz, T. J. Adv. Organomet. Chem. 1977, 16, 283. (f) Haines, R. J.; Leigh, G.J. Chem. Soc. Rev. 1975, 4, 1. (g) Mol, J. C.; Moulijn, J. A. Adv. Catal. 1975, 24, 131. (h) Dall'Asta, G. Rubber Chem. Techn. 1974, 47, 511.
- (4) Masuda, T.; Sasaki, N.; Higashimura, T. Macromolecules 1975, 8, 717.
- (5) Katz, T. J.; Lee, S. J. J. Am. Chem. Soc. 1980, 102, 422.

(6) (a) Katz, T. J. ; Lee, S. J. ; Nair, M.; Savage, E. B. Ibid. 1980, 102, 7940. (b) Katz, T. J.; Savage, E. B.; Lee, S. J.; Nair, M. Ibid. 1980, 102, 7942.

(7) Tungsten hexachloride alone does not metathesize olefins that are unstrained,⁸ but it does metathesize examples that are strained (like norbornene and dicyclopentadiene).¹⁰ Küpper demonstrated that combined with more reactive olefins it would induce the reactions of those that are less reactive,¹¹ and Makovetskii et al. very recently discovered results similar to some of those reported here, in which it is effective in combination with acetylenes.¹² Phenylacetylene is also recorded among additives that enhance the reactivity of tungsten hexachloride plus tetrabutyltin.¹³

(8) In the presence of oxygen, but not in its absence, WCl_6 is reported to polymerize cyclopentene to polypentenamer.⁹ Similar, but not identical, experiments (see Table I) failed for us. After a long time (22 h) at 30 °C with WCl_6 in air in the absence of solvent, cyclopentene gave a 2.6% yield of polymer that was 80% saturated (1H NMR analysis). In an equal volume of chlorobenzene (not toluene and not distilled from triethylaluminum) after the same amount of time it did not polymerize appreciably.

(9) Amass, A.J.; McGourtey, T.A. Eur. Polym. J. 1980, 16, 235.

(10) Oshika, T.; Tabushi, H. Bull. Chem. Soc. Jpn. 1968, 41, 211.

(11) Küpper, F.-W. Angew. Makromol. Chem. 1979, 80, 207.

(12) Makovetskii, K. L.; Red'kina, L. I.; Oreshkin, I. A. Izv. Akad.

Nauk SSSR, Ser. Khim. 1981, 1928.

(13) Ichikawa, K.; Watanabe, O.; Takagi, T.; Fukuzumi, K. J. Catal.
1976, 44, 416.

(14) 99.6% *cis*, containing 0.25% n-pentane.

(15) The volume of chlorobenzene equalled that of the cis-2-pentene.

No precaution was taken to exclude air.

(16) Katz, T. J.; Lee, S. J.; Acton, N. Tetrahedron Lett. 1976, 4247.

(17) The ^{13}C NMR spectra of polyalkenamers are described in reference 14, in the references in its footnote 18, and in (a) Carmen, C. J.; Wilkes, C.E. Macromolecules 1974, 7, 40. (b) Chen, H. Y. Appl. Polym. Spectrosc. 1978, 7.

(18) (a) Greco, A.; Pirinoli, F.; Dall'Asta, G. J. Organometal. Chem. 1973, 60, 115. (b) San Fillipo, Jr., J.; Sowinski, A. F.; Romano, L. J. J. Am. Chem. Soc. 1974, 97, 1599.

(19) Katz, T. J.; Hersh, W. H. Tetrahedron Lett. 1977, 585.

(20) See the results and summaries in references 3e, 6a, 16, and 19. Klupper's aluminum-free catalysts induce stereochemistries like those here.¹¹ For cyclopentene's metathesis decreasing amounts of aluminum halides decrease the stereoselectivity,²¹ and metals (like Sn²² and Si²³) whose halides are less acidic than aluminum's enhance stereoselectivity.

(21) (a) Günther, P. et al. Angew. Makromol. Chem. 1971, 14, 87 and 16/17, 27. (b) Ivin, K. J.; Laverty, D. T.; Rooney, J. J. Makromol. Chem. 1977, 178, 1545.

(22) Ivin, K. J.; Lapienis, G.; Rooney, J. J. Polymer 1980, 21, 367.

(23) Oreshkin, I. A. et al. Eur. Polym. J. 1977, 13, 447.

Table I. Yields, Stereochemistries, and Molecular Weights of Polymers Prepared According to Equation 2 When $x = 1, 4$, or 0 .^a

cyclo- alkene ^b	Reaction			Yield (%) ^c	% cis		$\bar{M}_w x$ 10^{-3}	$\bar{M}_n x$ 10^{-3}
	tempera- ture (°C)	time	x		IR ^d	^{13}C NMR ^e		
5	24	320 s	1	9.8±1	61.1±2	40.1±2	155	89
5	24	320 s	0	0.065±0.02				
8	30	5 h	4	21.4±0.4	79.4±0.7	74.5±0.6		
8	30	5 h	0	0.3±0				
8	30	19 h	1	17.2±1	86.1±1	85.0±0.9	223	127
8	30	19 h	0	0.9±0.7				
7	30	24 h	1	57.6	73.3±0.7	67.9±2		
7	30	24 h	0	0.8				

^a Cyclopentene and cyclooctene were diluted with equal volumes of chlorobenzene, but cycloheptene was not. Reactions were effected in the atmosphere. ^bThe number of carbons in the cycloalkene.

^cAfter dissolving in methylene chloride, precipitating with methanol, washing with aqueous ammonia and methanol (not for cycloheptene), and vacuum drying. The average deviation listed is that of three experiments with cyclopentene and two with cyclooctene. ^dMeasured by the IR spectra of thin films as described in footnote 16 of reference 16. The average deviations are of 2-4 measurements. ^eMeasured by the intensities of resonances near 32 and 27 ppm. Spectra were

acquired using CDCl_3 solutions, a 20 MHz spectrometer with proton noise decoupling and no relaxation delay (see note 17). ^fMeasured by gel-permeation chromatography using tetrahydrofuran solutions, a refractive index monitor, and 5 Waters Associates μ-styragel columns (10^6 , 10^5 , 10^4 , 10^3 , and 500 \AA). The curves are bimodal (ref 6b), and it is the analyses of the high-molecular weight peaks that are recorded. The values recorded are half the weights of the polystyrenes that would exhibit the chromatograms observed (see Table II in ref 6a and Table I in ref 6b). ^gNot measured, but $[\eta] = 6.15 \pm 0.2 \text{ dL/g}$ in toluene at 29.7°C .

Table II. The Dependence of the Yields and Molecular Weights of Polypentenamers Prepared According to Equation 2 (after 8 min at 21 °C) on the Amount of Phenylacetylene in the Reaction Mixture.^a

x^b	$\bar{M}_n \cdot 10^{-3} \cdot e^c$	Yield (%) ^{d,e}
0	0.12	33.5±7
0.31	3.6±3	143.6±5
1.25	5.6±4	16.8±4
2.5	14.9±3	79.1±0.2
3.75	(4.8)	22.5±0.3
7.5	(4.2)	15.2±0.02
10		

^aCyclopentene was diluted with an equal volume of chlorobenzene. Reactions were effected in the atmosphere.

^bThe number of moles of phenylacetylene per 200 moles of cyclopentene. ^c $\bar{M}_n \cdot 10^{-3}$ (weight of polymer/weight of cyclopentene + phenylacetylene). If the yield of polyphenylacetylene were 75% the yields of polypentenamer would (for $x \leq 3.75$) be 0.12, 3.4, 5.0, 15.8, and 13.4. For $x = 7.5$ and 10 however, much of the polymer is polyphenylacetylene, and if its yield were 75%, that of polypentenamer would be negligible.

^dWorked up as in Table I, note c. ^eSee footnote f in Table I. The averages and standard deviations of measurements made using a refractive index monitor and a detector of light absorption at 400 nm. The figures on the last two lines were estimated by supposing that the single peaks observed could be dissected into two overlapping components.

TECHNICAL REPORT DISTRIBUTION LIST, GEN

<u>No.</u> <u>Copies</u>		<u>No.</u> <u>Copies</u>
2	Office of Naval Research Attn: Code 472 800 North Quincy Street Arlington, Virginia 22217	U.S. Army Research Office Attn: CRD-AA-IP P.O. Box 1211 Research Triangle Park, N.C. 27709
1	ONR Branch Office Attn: Dr. George Sandoz 536 S. Clark Street Chicago, Illinois 60605	Naval Ocean Systems Center Attn: Mr. Joe McCartney San Diego, California 92152
1	ONR Area Office Attn: Scientific Dept. 715 Broadway New York, New York 10003	Naval Weapons Center Attn: Dr. A. B. Amster, Chemistry Division China Lake, California 93555
1	ONR Western Regional Office 1030 East Green Street Pasadena, California 91106	Naval Civil Engineering Laboratory Attn: Dr. R. W. Drisko Port Hueneme, California 93401
1	ONR Eastern/Central Regional Office Attn: Dr. L. H. Peebles Building 114, Section D 666 Summer Street Boston, Massachusetts 02210	Department of Physics & Chemistry Naval Postgraduate School Monterey, California 93940
1	Director, Naval Research Laboratory Attn: Code 6100 Washington, D.C. 20390	Dr. A. L. Slafkosky Scientific Advisor Commandant of the Marine Corps (Code RD-1) Washington, D.C. 20380
1	The Assistant Secretary of the Navy (RE&S) Department of the Navy Room 4E736, Pentagon Washington, D.C. 20350	Office of Naval Research Attn: Dr. Richard S. Miller 800 N. Quincy Street Arlington, Virginia 22217
1	Commander, Naval Air Systems Command Attn: Code 310C (H. Rosenwasser) Department of the Navy Washington, D.C. 20360	Naval Ship Research and Development Center Attn: Dr. G. Bosmajian, Applied Chemistry Division Annapolis, Maryland 21401
12	Defense Technical Information Center Building 5, Cameron Station Alexandria, Virginia 22314	Naval Ocean Systems Center Attn: Dr. S. Yamamoto, Marine Sciences Division San Diego, California 91232
1	Dr. Fred Saalfeld Chemistry Division, Code 6100 Naval Research Laboratory Washington, D.C. 20375	Mr. John Boyle Materials Branch Naval Ship Engineering Center Philadelphia, Pennsylvania 19112

TECHNICAL REPORT DISTRIBUTION LIST, GENNo.
Copies

Dr. Rudolph J. Marcus
Office of Naval Research
Scientific Liaison Group
American Embassy
APO San Francisco 96503 1

Mr. James Kelley
DTNSRDC Code 2803
Annapolis, Maryland 21402 1

Mr. A. M. Anzalone
Administrative Librarian
PLASTEC/ARRADCOM
Bldg 3401
Dover, New Jersey 07801 1

TECHNICAL REPORT DISTRIBUTION LIST, 356B

<u>No.</u>	<u>Copies</u>	<u>No.</u>	<u>Copies</u>
Dr. C. L. Shilling Union Carbide Corporation Chemical and Plastics Tarrytown Technical Center Tarrytown, New York	1	Dr. E. Fischer, Code 2853 Naval Ship Research and Development Center Annapolis Division Annapolis, Maryland 21402	1
Dr. R. Soulen Contract Research Department Pennwalt Corporation 900 First Avenue King of Prussia, Pennsylvania 19406	1	Dr. Martin H. Kaufman, Head Materials Research Branch (Code 4542) Naval Weapons Center China Lake, California 93555	1
Dr. A. G. MacDiarmid University of Pennsylvania Department of Chemistry Philadelphia, Pennsylvania 19174	1	Dr. C. Allen University of Vermont Department of Chemistry Burlington, Vermont 05401	1
Dr. H. Allcock Pennsylvania State University Department of Chemistry University Park, Pennsylvania 16802	1	Professor R. Drago Department of Chemistry University of Illinois Urbana, Illinois 61801	1
Dr. M. Kenney Case-Western University Department of Chemistry Cleveland, Ohio 44106	1	Professor H. A. Titus Department of Electrical Engineering Naval Postgraduate School Monterey, California 93940	1
Dr. R. Lenz University of Massachusetts Department of Chemistry Amherst, Massachusetts 01002	1	COL R. W. Bowles, Code 100M Office of Naval Research 800 N. Quincy Street Arlington, Virginia 22217	1
Dr. M. David Curtis University of Michigan Department of Chemistry Ann Arbor, Michigan 48105	1	Professor T. Katz Department of Chemistry Columbia University New York, New York 10027	1
NASA-Lewis Research Center Attn: Dr. T. T. Serafini, MS 49-1 21000 Brookpark Road Cleveland, Ohio 44135	1	Professor James Chien Department of Chemistry University of Massachusetts Amherst, Massachusetts 01002	1
Dr. J. Griffith Naval Research Laboratory Chemistry Section, Code 6120 Washington, D.C. 20375	1	Professor Malcolm B. Polk Department of Chemistry Atlanta University Atlanta, Georgia 30314	1
Dr. G. Goodman Globe-Union Incorporated 5757 North Green Bay Avenue Milwaukee, Wisconsin 53201	1	Dr. G. Bryan Street IBM Research Laboratory San Jose, California 95193	1

7-8